

Extra Strength Excedrin®

Product Information

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg
Aspirin 250 mg
Caffeine 65 mg

Purposes

Pain reliever
Pain reliever
Pain reliever aid

Uses

• temporarily relieves minor aches and pains due to:

- headache
 - a cold
 - arthritis
 - muscular aches
 - sinusitis
 - toothache
 - premenstrual & menstrual cramps
-

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen and aspirin or other pain relievers/fever reducers. Acetaminophen and aspirin may cause liver damage and stomach bleeding.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - with any other products containing acetaminophen. Taking more than directed may cause liver damage.
-

Ask a doctor before use if you have

- asthma
 - ulcers
 - bleeding problems
 - stomach problems such as heartburn, upset stomach, or stomach pain that do not go away or return
-

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- anticoagulation (thinning of the blood)
 - diabetes
 - gout
 - arthritis
-

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.

- new symptoms occur
 - symptoms do not get better or worsen
 - ringing in the ears or loss of hearing occurs
 - painful area is red or swollen
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
-

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
 - drink a full glass of water with each dose
 - adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
 - children under 12 years: ask a doctor
-

Other information

- store at room temperature
 - read all product information before using. Keep the box for important information.
-

Inactive ingredients

benzoic acid, carnauba wax, FD&C blue #1*, hydroxypropylcellulose, hypromellose, microcrystalline cellulose, mineral oil, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide*

*may contain these ingredients

Questions or comments? 1-800-468-7746