

# Excedrin Extra Strength Caplets (24ct)

## Drug Facts

---

### **Active ingredients (in each caplet)**

### **Purposes**

Acetaminophen 250 mg ..... Pain reliever

Aspirin 250 mg (NSAID\*) ..... Pain reliever

Caffeine 65 mg ..... Pain reliever aid

\*nonsteroidal anti-inflammatory drug

---

### **Uses**

Temporarily relieves minor aches and pains due to:

- headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps
- 

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a

doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

---

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
  - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- 

### **Ask a doctor before use if**

- you have liver disease
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
  - you are taking a diuretic
  - you have asthma
-

### Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
  - any other drug or are under a doctor's care for any serious condition
- 

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

---

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

---

### ***Directions***

- do not use more than directed
  - drink a full glass of water with each dose
  - adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
  - children under 12 years: ask a doctor
- 

### ***Other information***

- store at 20°-25°C (68°-77°F)
  - close cap tightly after use
  - read all product information before using. Keep this box for important information.
- 

### ***Inactive ingredients***

benzoic acid, carnauba wax, FD&C blue no. 1, hypromellose, light mineral oil, low-substituted hydroxypropyl cellulose, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

---

# Excedrin Extra Strength Caplets (100ct, 200ct & 300ct)

## Drug Facts

---

### **Active ingredients (in each caplet)**

### **Purposes**

Acetaminophen 250 mg ..... Pain reliever

Aspirin 250 mg (NSAID\*) ..... Pain reliever

Caffeine 65 mg ..... Pain reliever aid

\*nonsteroidal anti-inflammatory drug

---

### **Uses**

Temporarily relieves minor aches and pains due to:

- headache
  - a cold
  - arthritis
  - muscular aches
  - toothache
  - premenstrual and menstrual cramps
- 

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a

doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

---

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
  - with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- 

### **Ask a doctor before use if**

- you have liver disease
  - stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
  - you are taking a diuretic
  - you have asthma
-

### Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
  - any other drug or are under a doctor's care for any serious condition
- 

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition.

---

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

---

### ***Directions***

- do not use more than directed
  - drink a full glass of water with each dose
  - adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
  - children under 12 years: ask a doctor
- 

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
  - close cap tightly after use
  - read all product information before using. Keep this box for important information.
- 

### ***Inactive ingredients***

corn starch, crospovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

---